

## Recommended vaccinations for children up to 6 years old

If your child misses a shot, you don't need to start over, just go back to your child's doctor for the next shot. Talk with your child's doctor if you have any questions.

| Age            | Vaccine & Dose                                                                                                                                                                     | Age                | Vaccine & Dose                                                                                                                                |
|----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|-----------------------------------------------------------------------------------------------------------------------------------------------|
| At birth       | <b>Hepatitis B</b> (Dose 1 of 3)                                                                                                                                                   | 6 months or older  | <b>Influenza</b> (Dose 1 of 2)<br><b>Influenza</b> (Dose 2 of 2 a month later and one dose yearly thereafter)                                 |
| 1 to 2 months  | <b>Hepatitis B</b> (Dose 2 of 3)                                                                                                                                                   | 12 to 15 months    | <b>Hib</b> (Dose 4 of 4)<br><b>Pneumococcal conjugate</b> (PCV13) (Dose 4 of 4)<br><b>MMR</b> (Dose 1 of 2)<br><b>Varicella</b> (Dose 1 of 2) |
| 2 months       | <b>DTaP</b> (Dose 1 of 5)<br><b>Hib</b> (Dose 1 of 4)<br><b>Polio</b> (IPV) (Dose 1 of 4)<br><b>Pneumococcal conjugate</b> (PCV13) (Dose 1 of 4)<br><b>Rotavirus</b> (Dose 1 of 3) | 12 to 23 months    | <b>Hepatitis A</b> (Dose 1 of 2)                                                                                                              |
| 4 months       | <b>DTaP</b> (Dose 2 of 5)<br><b>Hib</b> (Dose 2 of 4)<br><b>Polio</b> (IPV) (Dose 2 of 4)<br><b>Pneumococcal conjugate</b> (PCV13) (Dose 2 of 4)<br><b>Rotavirus</b> (Dose 2 of 3) | 15 to 18 months    | <b>DTaP</b> (Dose 4 of 5)                                                                                                                     |
| 6 months       | <b>DTaP</b> (Dose 3 of 5)<br><b>Hib</b> (Dose 3 of 4)<br><b>Pneumococcal conjugate</b> (PCV13) (Dose 3 of 4)<br><b>Rotavirus</b> (Dose 3 of 3)                                     | 18 months or older | <b>Hepatitis A</b> (Dose 2 of 2, follows 6 months after Dose 1)                                                                               |
| 6 to 18 months | <b>Hepatitis B</b> (Dose 3 of 3)<br><b>Polio</b> (IPV) (Dose 3 of 4)                                                                                                               | 4 - 6 years        | <b>DTaP</b> (Dose 5 of 5)<br><b>Polio</b> (IPV) (Dose 4 of 4)<br><b>MMR</b> (Dose 2 of 2)<br><b>Varicella</b> (Dose 2 of 2)                   |

NOTE: Consult your child's pediatrician or health-care professional about catch-up vaccinations and vaccines for travelers or if your child has special health conditions.

SOURCE: U.S. Department of Health and Human Services